



162/345



a-c, e (independently selected) = 0 or 1;
 d = 0;
 R = modifying group, mannose, oligo-
 mannose.

FIG. 39A

163/345




FIG. 39B




FIG. 39C

164/345




FIG. 39D




FIG. 39E

165/345




FIG. 39F







FIG. 39G

166/345

2 peptides
 A and A' - N-linked sites
 B - O-linked sites



Alternate structure
 for some N-linked
 structures of A.

a-d, i, n-u (independently selected) = 0 or 1.
 aa, bb (independently selected) = 0 or 1.
 e-h (independently selected) = 0 to 6.
 j-m (independently selected) = 0 to 20.
 v-z = 0; R = polymer, glycoconjugate.

FIG. 40A

167/345

CHO, BHK, 293s cells, Vero, MDCK, HEKC expressed
Factor VIII.

e-h = 1 to 4;

aa, bb, a-d, j-m, i, n-u (independently selected) = 0 or 1;

v-z = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG, ST3Gal3

e-h = 1 to 4;

aa, bb, a-d, i, n, q-u (independently selected) = 0 or 1;

o, p, z = 0; j-m, v-y (independently selected) = 0 or 1;

R = PEG.

FIG. 40B

CHO, BHK, 293S cells, Vero, MDCK, 293S, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, a-d, j-m, i, n-u (independently selected) = 0 or 1;

v-z = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG, ST3Gal3
 3. ST3Gal1, CMP-SA

e-h = 1 to 4;

aa, bb, a-d, i, n, p-u (independently selected) = 0 or 1;

o, z = 0; j-m, v-y (independently selected) = 0 or 1;

R = PEG.

FIG. 40C

168/345

CHO, BHK, 293s cells, Vero, MDCK, HEKC
expressed Factor VIII.
e-h = 1 to 4;
aa, bb, a-d, j-m, i, n-u (independently selected)=0 or 1;
v-z = 0.

↓
1. CMP-SA-PEG, ST3Gal3

e-h = 1 to 4;
aa, bb, a-d, i, n-u (independently selected) = 0 or 1;
z = 0; j-m, v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 40D

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.
e-h = 1 to 4;
aa, bb, a-d, j-m, i, n-u (independently selected) 0 or 1;
v-z = 0.

↓
1. CMP-SA-PEG, ST3Gal1

e-h = 1 to 4;
aa, bb, a-d, i, n-u (independently selected) = 0 or 1;
z = 0; j-m, v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 40E

169/345

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, a-d, j-m, i, n-u (independently selected)=0 or 1;

v-z = 0.



1. CMP-SA-PEG, α 2,8-ST

e-h = 1 to 4;

aa, bb, a-d, i, n-y (independently selected) = 0 or 1;

z = 0; j-m (independently selected) = 0 to 2;

v-y (independently selected) = 1,

when j-m (independently selected) is 2;

R = PEG.




FIG. 40F

170/345

2 peptides

A or A' - N-linked sites

B - O-linked sites



a-d, i, n-u, (independently selected) = 0 or 1.

aa, bb, cc, dd (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 6.

j-m (independently selected) = 0 to 20.

$$y-z = 0;$$

R = modifying group, mannose, oligo-mannose.

R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 40G

171/345

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, cc, a-d, j-m, i, n-u (independently selected) = 0 or 1;

dd, v-z = 0.

↓ 1. CMP-SA-levulinate, ST3Gal3,
2. H₄N₂-PEG

e-h = 1 to 4;

aa, bb, cc, a-d, i, n-u (independently selected) = 0 or 1;

dd, z = 0; j-m, v-y (independently selected) = 0 or 1;

R = PEG.

FIG. 40H

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, cc, a-d, j-m, i, n-u (independently selected) = 0 or 1;

dd, v-z = 0.

↓ 1. endo-H
2. galactosyltransferase, UDP-Gal-PEG

e-h = 1 to 4;

aa, bb, dd, a-d, i, j-u (independently selected) = 0 or 1;

cc, v-z = 0; R' = -Gal-PEG.

FIG. 40I

172/345

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, cc, a-d, j-m, i, n-u (independently selected) = 0 or 1;

dd, v-z = 0.

- ↓
1. ST3Gal3, CMP-SA
 2. endo-H
 3. galactosyltransferase, UDP-Gal-PEG

e-h = 1 to 4;

aa, bb, dd, a-d, i, j-u (independently selected) = 0 or 1;

cc, v-z = 0; R' = -Gal-PEG.

FIG. 40J

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, cc, a-d, j-m, i, n-u (independently selected) = 0 or 1;

dd, v-z = 0.

- ↓
1. mannosidases
 2. GNT 1 & 2, UDP-GlcNAc
 3. galactosyltransferase, UDP-Gal-PEG

e-h = 1 to 4;

aa, a-d, i, j-y (independently selected) = 0 or 1;

bb, cc, dd, z = 0; R = PEG.

FIG. 40K

173/345

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, cc, a-d, j-m, i, n-u (independently selected) = 0 or 1;

dd, v-z = 0.

1. mannosidases

2. GNT-1, 2, 4 & 5; UDP-GlcNAc

↓ 3. galactosyltransferase, UDP-Gal

4. ST3Gal3, CMP-SA

e-h = 1 to 4;

aa, bb, cc, a-d, i, j-q (independently selected) = 0 or 1;

dd, v-z = 0.

FIG. 40L

CHO, BHK, 293S cells, Vero, MDCK, HEKC
expressed Factor VIII.

e-h = 1 to 4;

aa, bb, cc, a-d, j-m, i, n-u (independently selected) = 0 or 1;

dd, v-z = 0.

1. mannosidases

↓ 2. GNT-1, UDP-GlcNAc-PEG


e-h = 0 to 4;

aa, a-d, i, j-y (independently selected) = 0 or 1;

bb, cc, dd, z = 0.

FIG. 40M

174/345



a-d, i, q-u (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 6.

j-m (independently selected) = 0 to 100.

v-y = 0; R = polymer.

FIG. 41A

175/345

CHO, BHK, 293 cells, Vero expressed Urokinase.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG (16 mol eq),
ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 1,
when j-m (independently selected) is 1;
R = PEG.

FIG. 41B

CHO, BHK, 293 cells, Vero expressed Urokinase.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG (1.2 mol eq),
ST3Gal3
 3. CMP-SA (16 mol eq), ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 41C

176/345

NSO expressed Urokinase.
 a-d, i-m, q-u (independently selected) = 0 or 1;
 e-h = 1; v-y = 0;
 Sia (independently selected) = Sia or Gal.

- ↓
1. Sialidase and α -galactosidase
 2. α -Galactosyltransferase, UDP-Gal
 - ↓ 3. CMP-SA-PEG, ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
 e-h = 1; v-y (independently selected) = 1,
 when j-m (independently selected) is 1;
 R = PEG.

FIG. 41D

CHO, BHK, 293 cells, Vero expressed Urokinase.
 a-d, i-m, q-u (independently selected) = 0 or 1;
 e-h = 1; v-y = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG (16 mol eq),
ST3Gal3
 - ↓ 3. CMP-SA, ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
 e-h = 1; v-y (independently selected) = 0 or 1;
 R = PEG.

FIG. 41E

177/345

CHO, BHK, 293 cells, Vero expressed Urokinase.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. CMP-SA-levulinate, ST3Gal3,
buffer, salt
 2. H_4N_2 -PEG

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 41F


CHO, BHK, 293 cells, Vero expressed Urokinase.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. CMP-SA, $\alpha 2,8$ -ST

a-d, i, q-u (independently selected) = 0 or 1;
e-h = 1;
j-m (independently selected) = 0-20;
v-y (independently selected) = 0.

FIG. 41G

178/345



a-d, i, n, p-u (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 6.

j-m (independently selected) = 0 to 100.

v-y = 0;

R = modifying group, mannose, oligo-mannose;

R' = H, glycosyl residue, modifying group,
glycoconjugate.

FIG. 41H

179/345




FIG. 41I




FIG. 41J

180/345

CHO, BHK, 293 cells, Vero expressed Urokinase.

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; n, v-y = 0.

- ↓
1. CMP-SA-linker-SA-CMP, ST3Gal3
 2. ST3Gal1, desialylated Urokinase produced in CHO.
 - ↓ 3. CMP-SA, ST3Gal3, ST3Gal1

a-m, q-u (independently selected) = 0 or 1;

p = 1; n = 0;

v-y (independently selected) = 0 or 1;

R = linker-Urokinase.

FIG. 41K

Isolated Urokinase.

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; v-y = 0; n = 0;

Sia (independently selected) = Sia or SO₄;

Gal (independently selected) = Gal or GalNAc;

GlcNAc (independently selected) = GlcNAc or GlcNAc-Fuc.

- ↓
1. sulfohydrolase
 2. CMP-SA-PEG, sialyltransferase

a-d, i-m, q-u (independently selected) = 0 or 1;

n = 0; e-h = 1; Sia = Sia;

Gal (independently selected) = Gal or GalNAc;

GlcNAc (independently selected) = GlcNAc or GlcNAc-Fuc.

v-y (independently selected) = 0 or 1;

R = PEG.

FIG. 41L

181/345

Isolated Urokinase.

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; n = 0; v-y = 0;

Sia (independently selected) = Sia or SO₄;

Gal (independently selected) = Gal or GalNAc;

GlcNAc (independently selected) = GlcNAc or GlcNAc-Fuc.



1. sulfohydrolase, hexosaminidase

2. UDP-Gal-PEG, galactosyltransferase

a-d, i, q-u (independently selected) = 0 or 1;



e-h = 1; j-n = 0; Gal (independently selected) = Gal;

GlcNAc (independently selected) = GlcNAc or GlcNAc-Fuc;

v-y (independently selected) = 0 or 1; R = PEG.

FIG. 41M

182/345



a-d, i, q-u (independently selected) = 0 or 1.
 e-h (independently selected) = 0 to 6.
 j-m (independently selected) = 0 to 100.
 v-y = 0; R = polymer, glycoconjugate.

FIG. 42A

183/345

CHO, BHK, 293 cells, Vero expressed DNase I.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG (16 mol eq),
ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1;
v-y (independently selected) = 1,
when j-m (independently selected) is 1;
R = PEG.

FIG. 42B

CHO, BHK, 293 cells, Vero expressed DNase I.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG (1.2 mol eq), ST3Gal3
 3. CMP-SA (16 mol eq), ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 42C

184/345

NSO expressed DNase I.

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; v-y = 0;

Sia (independently selected) = Sia or Gal.

- ↓
1. Sialidase and α -galactosidase
 2. α -Galactosyltransferase, UDP-Gal
 3. CMP-SA-PEG, ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; v-y (independently selected) = 1,

when j-m (independently selected) is 1;

R = PEG.

FIG. 42D

CHO, BHK, 293 cells, Vero expressed DNase I.

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; v-y = 0.

- ↓
1. Sialidase
 2. CMP-SA-PEG (16 mol eq), ST3Gal3
 3. CMP-SA, ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;

e-h = 1; v-y (independently selected) = 0 or 1;

R = PEG.

FIG. 42E

185/345

CHO, BHK, 293 cells, Vero expressed DNase I.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. CMP-SA-levulinate, ST3Gal3,
buffer, salt
 2. H₄N₂-PEG

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 42F

CHO, BHK, 293 cells, Vero expressed DNase I.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

- ↓
1. CMP-SA, α 2,8-ST

a-d, i, q-u (independently selected) = 0 or 1;
e-h = 1;
j-m (independently selected) = 0-20;
v-y (independently selected) = 0.


FIG. 42G

187/345

Insect cell expressed DNase I.

a-d, f, h, j-n, s, u, v-y' = 0;

e, g, i, q, r, t (independently selected) = 0 or 1.



1. GNT's 1,2,4,5, UDP-GlcNAc

2. Galactosyltransferase, UDP-Gal-PEG

a-i, q-u (independently selected) = 0 or 1; j-n = 0;

v-y (independently selected) = 1,

when e-h (independently selected) is 1;

R = PEG.

FIG. 42I

Yeast expressed DNase I.

a-n = 0;

q-y (independently selected) = 0 to 1;

p = 1; R (branched or linear) = Man, oligomannose.



1. Endoglycanase

2. Galactosyltransferase, UDP-Gal

3. CMP-SA-PEG, ST3Gal3

a-n, p-y = 0; n (independently selected) = 0 or 1;

R' = -Gal-Sia-PEG.

FIG. 42J

188/345


CHO, BHK, 293 cells, Vero expressed DNase I.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; n, v-y = 0.

- ↓
1. CMP-SA-linker-SA-CMP, ST3Gal3
 2. ST3Gal1, desialylated alpha-1-Proteinase inhibitor.
 3. CMP-SA, ST3Gal3, ST3Gal1

a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker- alpha-1-Proteinase inhibitor.

FIG. 42K

189/345



a-d, i, r-u (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 4.

j-m (independently selected) = 0 or 1.

n, v-y = 0; z = 0 or 1;

R = modifying group, mannose, oligo-mannose;

R' = H, glycosyl residue, modifying group,
glycoconjugate.

FIG. 43A

190/345




FIG. 43B




FIG. 43C

191/345

Yeast expressed Insulin.

a-n = 0; r-y (independently selected) = 0 to 1;
z = 1;


R (branched or linear) = Man, oligomannose or
polysaccharide.

- ↓
1. Endo-H
2. Galactosyltransferase, UDP-Gal-PEG

a-m, r-z = 0; n = 1; R' = -Gal-PEG.

FIG. 43D

192/345



a-c, e (independently selected) = 0 or 1;
 d = 0; R = polymer

FIG. 43E

193/345




FIG. 43F




FIG. 43G

194/345









FIG. 43H

195/345



a-c, e (independently selected) = 0 or 1;
 d = 0; R = modifying group, mannose,
 oligo-mannose.

FIG. 43I

196/345




FIG. 43J




FIG. 43K

197/345


E. coli expressed Insulin (N)—no mucin peptide.
a-e, n = 0.

- ↓
1. NHS-CO-linker-SA-CMP
 2. ST3Gal3, asialo-transferrin
 3. CMP-SA, ST3Gal3

a-e = 0; n = 1;
R' = linker-transferrin.

FIG. 43L

198/345



a-d, i, n-u, aa (independently selected) = 0 or 1.
 e-h (independently selected) = 0 to 6.
 j-m (independently selected) = 0 to 100.
 v-y = 0; R = polymer, glycoconjugate.

FIG. 44A

199/345

CHO, BHK, 293 cells, Vero expressed M-antigen.
a-d, i-m, o-u, aa (independently selected) = 0 or 1;
n, e-h = 1; v-z = 0.

- ↓
1. Sialidase
 2. CMP-SA-linker-lipid-A,
ST3Gal3

a-d, i-m, q-u, aa (independently selected) = 0 or 1;
o, p, z = 0; n, e-h = 1;
v-y (independently selected) = 1,
when j-m (independently selected) is 1;
R = linker-lipid-A.

FIG. 44B

CHO, BHK, 293 cells, Vero expressed M-antigen.
a-d, i-m, o-u, aa (independently selected) = 0 or 1;
n, e-h = 1; v-z = 0.

- ↓
1. sialidase
 2. CMP-SA-linker-tetanus toxin, ST3Gal1
 3. CMP-SA, ST3Gal3

a-d, i-m, p-u, z, aa (independently selected) = 0 or 1;
o, v-y = 0; n, e-h = 1; R = tetanus toxin.

FIG. 44C

200/345

NSO expressed M-antigen.

a-d, i-n, o-u, aa (independently selected) = 0 or 1;

e-h = 1; v-z = 0;

Sia (independently selected) = Sia or Gal.

- ↓
1. α -galactosidase
 2. CMP-SA, ST3Gal3
 2. CMP-SA-KLH, ST3Gal1

a-d, i-n, p-u, z, aa (independently selected) = 0 or 1;

e-h = 1; o, v-y = 0;

z = 1, when p = 1;

R = KLH.

FIG. 44D

Yeast expressed M-antigen.

a-p, z = 0; q-y, aa (independently selected) = 0 to 1;

R (branched or linear) = Man, oligomannose;

GalNAc = Man.

- ↓
1. α 1,2-mannosidase
 2. GNT 1,
- ↓ UDP-GlcNAc-linker-diphtheria toxin.

e, q, l, m, r, t, u, v, aa (independently selected) = 0 or 1;

a-d, f-h, j, k, n-p, s, w-z = 0;

Sia = Man; R = linker-diphtheria toxin.

FIG. 44E

201/345




FIG. 44F





FIG. 44G

202/345



a-d, i, n, q-u, aa, bb, (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 6.

j-p (independently selected) = 0 to 100.

Cc, v-y = 0;

R = modifying group, mannose, oligo-mannose.

R' = H, glycosyl residue, modifying group,
glycoconjugate.

FIG. 44H

203/345




FIG. 44I




FIG. 44J

204/345

Yeast expressed M-antigen.

a-p, z, cc = 0;

q-y, aa (independently selected) = 0 to 1; bb = 1;

R (branched or linear) = Man, oligomannose;

GalNAc = Man.

1. mannosidases

2. GNT 1 & 2, UDP-GlcNAc

3. UDP-Gal, Galactosyltransferase,


4. CMP-SA, sialyltransferase

a, c, e, g, j, l, q, r, t, aa (independently selected) = 0 or 1;

b, d, f, h, k, m-p, s, u-z, cc = 0; bb = 1.

FIG. 44K

205/345



a-d, i, r-u (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 4.

j-m (independently selected) = 0 or 1.

n, v-y = 0; z = 0 or 1;

R = modifying group, mannose, oligo-mannose;

R' = H, glycosyl residue, modifying group,
glycoconjugate.

FIG. 45A

206/345




FIG. 45B




FIG. 45C

207/345





FIG. 45D

208/345



a-c, e (independently selected) = 0 or 1;
 d = 0;
 R = modifying group, mannose, oligo-
 mannose.

FIG. 45E

209/345




FIG. 45F




FIG. 45G

210/345




FIG. 45H




FIG. 45I

211/345




FIG. 45J





FIG. 45K

212/345



a-d, i-m, q-u, w, z, nn, ww, zz (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 4.

n, v-y = 0;

R = modifying group, mannose, oligo-mannose;

R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 46A

213/345

CHO, BHK, 293 cells, Vero or transgenic animals
expressed TNF Receptor IgG Fusion.
a-m, o-u, aa (independently selected) = 0 or 1;
n = 1; v-z = 0.

- ↓
1. CMP-SA, ST3Gal1
2. galactosyltransferase, UPD-Gal
3. CMP-SA-PEG, ST3Gal3

a-m, o-u, v-y, aa (independently selected) = 0 or 1;
n = 1; z = 0; R = PEG.

FIG. 46B

CHO, BHK, 293 cells, Vero expressed
TNF Receptor IgG Fusion.
a-m, o-u, aa (independently selected) = 0 or 1;
n = 1; v-z = 0.

- ↓
1. sialidase
2. CMP-SA-PEG, ST3Gal1

a-i, p-u, z, aa (independently selected) = 0 or 1;
n = 1; o, j-m, v-y = 0; R = PEG.

FIG. 46C

214/345

CHO, BHK, 293 cells, Vero expressed
TNF Receptor IgG Fusion.
a-m, o-u, aa (independently selected) = 0 or 1;
n = 1; v-z = 0.



1. galactosyltransferase, UPD-Gal-PEG

a-m, o-u, v-y, aa (independently selected) = 0 or 1;
n = 1; z = 0; R = PEG.

FIG. 46D

CHO, BHK, 293 cells, Vero or transgenic animals
expressed TNF Receptor IgG Fusion.
a-m, o-u, aa (independently selected) = 0 or 1;
n = 1; v-z = 0.



1. CMP-SA, ST3Gal1

2. galactosyltransferase, UPD-Gal-PEG

a-m, o-u, v-y, aa (independently selected) = 0 or 1;
n = 1; z = 0; R = PEG.

FIG. 46E

215/345

CHO, BHK, 293 cells, Vero or transgenic animals
expressed TNF Receptor IgG Fusion.
a-m, o-u, aa (independently selected) = 0 or 1;
n = 1; v-z = 0.

↓
1. CMP-SA-levulinate, ST3Gal1
2. H₄N₂-PEG

a-m, o-u, v-y, aa (independently selected) = 0 or 1;
n = 1; z = 0; R = PEG.

FIG. 46F


CHO, BHK, 293 cells, Vero expressed
TNF Receptor IgG Fusion.
a-m, o-u, aa (independently selected) = 0 or 1;
n = 1; v-z = 0.

↓
1. CMP-SA-PEG, α 2,8-ST

a-i, o, q-u, v-z, aa (independently selected) = 0 or 1;
n = 1; j-m, p (independently selected) = 0 to 2;
v-z (independently selected) = 1,
when j-m, p (independently selected) is 2;
R = PEG.

FIG. 46G

216/345



a-d, i, l, q-u (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 4.

j-k (independently selected) = 0 or 1.

M = 0 to 20.

n, v-y = 0; z = 0 or 1;

R = polymer, toxin, radioisotope-complex, drug, mannose, oligo-mannose.

R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 47A

217/345

CHO, BHK, 293 cells, Vero expressed Herceptin.
 a, c, i (independently selected) = 0 or 1;
 e, g, r, t = 1; b, d, f, h, j-m, n, s, u-y = 0;
 q, z = 1.



1. galactosyltransferase, UPD-Gal
2. CMP-SA-toxin, ST3Gal3

a, c, i, j, l (independently selected) = 0 or 1;
 e, g, r, t = 1; R = toxin;
 f, h, k, m, n, s, u-y = 0; q, z = 1;
 v-y (independently selected) = 51,
 when j, l (independently selected) is 1.

FIG. 47B

CHO, BHK, 293 cells, Vero or fungal expressed Herceptin.
 a, c, i (independently selected) = 0 or 1;
 e, g, r, t = 1; b, d, f, h, j-m, n, s, u-y = 0;
 q, z = 1.



1. galactosyltransferase,
UPD-Gal-Toxin

a, c, i (independently selected) = 0 or 1;
 e, g, r, t = 1; f, h, j-m, n, s, u-y = 0;
 q, z = 1; v-y (independently selected) = 1,
 when a, c (independently selected) is 1;
 R = toxin.

FIG. 47C

218/345

Fungi expressed Herceptin.

e, g, i, r, t (independently selected) = 0 or 1;

a-d, f, h, j-m, n, s, u-y = 0; q, z = 1.


- ↓ 1. Endo-H
2. Galactosyltransferase, UDP-Gal
3.. CMP-SA-radioisotope complex, ST3Gal3

a-m, r-z = 0; q, n = 1;

R' = -Gal-Sia-radioisotope complex.

FIG. 47D

219/345



a-d, i, p-u, (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 4.

j-m (independently selected) = 0 or 1.

n, v-y = 0; z = 0 or 1;

R = polymer, toxin, radioisotope-complex, drug, mannose, oligo-mannose.

R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 48A

220/345




FIG. 48B




FIG. 48C

221/345

Fungi expressed Synagis.

e, g, i, r, t (independently selected) = 0 or 1;


a-d, f, h, j-m, n, s, u-y = 0; q, z = 1.

- ↓
1. Endo-H
 2. Galactosyltransferase, UDP-Gal
 - 3.. CMP-SA-PEG, ST3Gal3

a-m, r-z= 0; q, n = 1; R' = -Gal-Sia-PEG.

FIG. 48D

222/345



a-d, i, q-u, w (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 6.

j-m (independently selected) = 0 to 20.

$$n, v-y=0; z=0 \text{ or } 1;$$

R = polymer, toxin, radioisotope-complex, drug, mannose, oligo-mannose.

R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 49A

223/345




FIG. 49B




FIG. 49C

224/345

Fungi expressed Remicade.


e, g, i, r, t (independently selected) = 0 or 1;
a-d, f, h, j-m, n, s, u-y = 0; q, z = 1.

- ↓
1. Endo-H
 2. Galactosyltransferase, UDP-Gal
 - 3.. CMP-SA-radioisotope complex, ST3Gal3

a-m, r-z = 0; q, n = 1;
R' = -Gal-Sia-radioisotope complex.

FIG. 49D

225/345



a-d, i, q-u (independently selected) = 0 or 1.

e-h (independently selected) = 0 to 4.

j-m (independently selected) = 0 or 1.

n, v-y = 0; z = 0 or 1;

R = modifying group, mannose, oligo-mannose;

R' = H, glycosyl residue, modifying group,
glycoconjugate.

FIG. 50A

226/345




FIG. 50B




FIG. 50C

227/345

Yeast expressed Reopro.

a-n = 0; r-y (independently selected) = 0 to 1;

z = 1;

R (branched or linear) = Man, oligomannose or polysaccharide.


1. Endo-H

2. Galactosyltransferase, UDP-Gal-PEG

a-m, r-z= 0; n = 1; R' = -Gal-PEG.

FIG. 50D

228/345



a-c, e (independently selected) = 0 or 1;
 d = 0; R = polymer

FIG. 50E

229/345




FIG. 50F




FIG. 50G

230/345






E. coli expressed Reopro-mucin fusion protein.
a-e = 0.

- ↓
1. GalNAc Transferase, UDP-GalNAc
 2. CMP-SA-PEG, sialyltransferase

c, d, e (independently selected) = 0 or 1;
a, b = 0; R = PEG.

FIG. 50H

231/345



a-c, e (independently selected) = 0 or 1;
d = 0; R = polymer, linker.

FIG. 50I

232/345




FIG. 50J




FIG. 50K

233/345

E. coli expressed Reopro(N)—no mucin peptide.
a-e, n = 0.

- ↓
1. NHS-CO-linker-SA-CMP
 2. ST3Gal3, asialo-transferrin
 3. CMP-SA, ST3Gal3

a-e = 0; n = 1; R' = linker-transferrin.

FIG. 50L